



US007488757B2

(12) **United States Patent**
Hoang et al.

(10) **Patent No.:** **US 7,488,757 B2**
(45) **Date of Patent:** **Feb. 10, 2009**

(54) **INVISIBLE ANTIMICROBIAL GLOVE AND
HAND ANTISEPTIC**

(75) Inventors: **Minh Quang Hoang**, Sandy, UT (US);
Donald Edward Hunt, Provo, UT (US)

(73) Assignee: **Becton, Dickinson and Company**,
Franklin Lakes, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 375 days.

(21) Appl. No.: **10/550,184**

(22) PCT Filed: **Mar. 24, 2004**

(86) PCT No.: **PCT/US2004/009009**

§ 371 (c)(1),
(2), (4) Date: **Jul. 24, 2006**

(87) PCT Pub. No.: **WO2004/084973**

PCT Pub. Date: **Oct. 7, 2004**

(65) **Prior Publication Data**

US 2006/0263323 A1 Nov. 23, 2006

Related U.S. Application Data

(60) Provisional application No. 60/457,054, filed on Mar.
24, 2003.

(51) **Int. Cl.**

A01N 25/00 (2006.01)
C11D 3/00 (2006.01)
A61K 31/045 (2006.01)
A61K 31/047 (2006.01)
A61K 31/05 (2006.01)
A61K 31/14 (2006.01)
A61K 31/155 (2006.01)
A61K 31/23 (2006.01)
A61K 31/235 (2006.01)

(52) **U.S. Cl.** **514/635**; 510/132; 514/483;
514/544; 514/545; 514/552; 514/636; 514/637;
514/642; 514/724; 514/738

(58) **Field of Classification Search** 424/404,
424/405; 510/132; 514/483, 545, 552, 635,
514/636, 637, 642, 724, 738

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,009,714 A 3/1977 Hammer
4,066,556 A 1/1978 Vaillancourt
4,294,594 A 10/1981 Sloane, Jr. et al.
4,309,992 A 1/1982 Dodak et al.
4,311,587 A 1/1982 Nose et al.
4,374,126 A 2/1983 Cardarelli et al.
4,416,772 A 11/1983 Sato et al.
4,560,382 A 12/1985 Isono
4,568,366 A 2/1986 Frederick et al.
4,678,606 A 7/1987 Akhter et al.

4,695,382 A 9/1987 Cronin
4,702,829 A 10/1987 Polaschegg et al.
4,714,556 A 12/1987 Ambrus et al.
4,738,668 A 4/1988 Bellotti et al.
4,787,974 A 11/1988 Ambrus et al.
4,861,485 A 8/1989 Fecondini
4,895,566 A 1/1990 Lee
5,053,339 A 10/1991 Patel
5,093,235 A 3/1992 Williams et al.
5,217,493 A 6/1993 Raad et al.
5,252,222 A 10/1993 Matkovich et al.
5,312,813 A 5/1994 Costerton et al.
5,366,505 A 11/1994 Farber
5,512,199 A 4/1996 Khan et al.
5,530,102 A 6/1996 Gristina et al.
5,554,147 A 9/1996 Batich et al.
5,556,760 A 9/1996 Nakamura et al.
5,638,812 A 6/1997 Turner
5,645,824 A 7/1997 Lim et al.
5,716,406 A 2/1998 Farber
5,723,132 A 3/1998 Tseng et al.
5,788,687 A 8/1998 Batich et al.
5,798,115 A 8/1998 Santerre et al.
5,817,063 A 10/1998 Turnbull
5,855,896 A 1/1999 Lim et al.
5,906,834 A 5/1999 Tseng
5,935,094 A 8/1999 Zupkas
6,017,334 A 1/2000 Rawls
6,022,551 A * 2/2000 Jampani et al. 424/405
6,022,748 A 2/2000 Charych et al.
6,048,337 A 4/2000 Svedman
6,080,423 A 6/2000 Charych et al.
6,106,287 A 8/2000 Yates
6,132,765 A 10/2000 DiCosmo et al.
6,159,007 A 12/2000 Sorensen
6,171,496 B1 1/2001 Patil
6,180,135 B1 1/2001 Charych et al.
6,180,584 B1 1/2001 Sawan et al.

(Continued)

Primary Examiner—John Pak
Assistant Examiner—Nathan W Schlientz
(74) *Attorney, Agent, or Firm*—The Webb Law Firm

(57) **ABSTRACT**


An alcohol based hand surgical scrub, which includes cat-
ionic anti-microbial agent preservatives, cationic polymer
film-forming agents and a skin emollient, and provides a long
term residual, anti-microbial “invisible glove” on the skin.
The composition provides an immediate reduction in bacter-
ial microbes. The polymer system creates an invisible film on
the skin following solvent evaporation. This invisible film
provides a lasting anti-microbial barrier on the skin which
acts to prevent microbial growth.

2 Claims, 1 Drawing Sheet

U.S. PATENT DOCUMENTS						
6,228,393	B1	5/2001	DiCosmo et al.	2003/0065292	A1 4/2003	Darouiche et al.
6,257,265	B1	7/2001	Brunner et al.	2003/0068667	A1 4/2003	Olson et al.
6,259,074	B1	7/2001	Brunner et al.	2003/0105143	A1 6/2003	Ammenodola et al.
6,277,652	B1	8/2001	Jo et al.	2003/0134783	A1 7/2003	Harshey et al.
6,282,444	B1	8/2001	Kroll et al.	2003/0153983	A1 8/2003	Miller et al.
6,306,422	B1	10/2001	Batich et al.	2003/0175812	A1 9/2003	Reppy et al.
6,306,598	B1	10/2001	Charych et al.	2003/0176367	A1 9/2003	Cali et al.
6,350,251	B1	2/2002	Prosl et al.	2003/0176848	A1 9/2003	Gibson et al.
6,391,001	B1	5/2002	Graham et al.	2003/0206875	A1 11/2003	Budny et al.
6,395,561	B1	5/2002	Charych et al.	2003/0215433	A1 11/2003	Kokai-Kun et al.
6,399,853	B1	6/2002	Roe et al.	2003/0224032	A1 12/2003	Read et al.
6,423,219	B1	7/2002	Chandler	2003/0229065	A1 12/2003	Levy et al.
6,428,491	B1	8/2002	Weiss	2004/0044299	A1 3/2004	Utsugi
6,436,885	B2	8/2002	Biedermann et al.	2004/0106912	A1 6/2004	Rosinskaya et al.
6,451,748	B1	9/2002	Taylor et al.	2004/0109852	A1 6/2004	Xu
6,468,649	B1	10/2002	Zhong	2004/0115721	A1 6/2004	Mao et al.
6,471,689	B1	10/2002	Joseph et al.	2004/0116845	A1 6/2004	Darouiche et al.
6,475,434	B1	11/2002	Darouiche	2004/0116896	A1 6/2004	Massengale
6,475,516	B2	11/2002	DiCosmo et al.	2004/0122511	A1 6/2004	Mangiardi et al.
6,479,727	B1	11/2002	Roe	2004/0126897	A1 7/2004	Prince et al.
6,501,002	B1	12/2002	Roe et al.	2004/0132164	A1 7/2004	Doyle et al.
6,540,915	B2	4/2003	Patil	2004/0132217	A1 7/2004	Prince et al.
6,562,295	B1	5/2003	Neuberger	2004/0172000	A1 9/2004	Roe et al.
6,579,539	B2	6/2003	Lawson et al.	2004/0180829	A1 9/2004	Bassler et al.
6,660,484	B2	12/2003	Charych et al.	2004/0188351	A1 9/2004	Thiele et al.
6,666,842	B1	12/2003	Sakai	2004/0230162	A1 11/2004	Tan
6,699,391	B2	3/2004	Baldrige et al.	2004/0235914	A1 11/2004	Ammendola et al.
6,713,660	B1	3/2004	Roe et al.	2004/0249441	A1 12/2004	Miller et al.
6,719,907	B2	4/2004	Collins et al.	2004/0254545	A1 12/2004	Rider et al.
6,719,991	B2	4/2004	Darouiche et al.	2005/0008671	A1 1/2005	Antwerp
6,723,350	B2	4/2004	Burrell et al.	2005/0010158	A1 1/2005	Brugger et al.
6,723,689	B1*	4/2004	Hoang et al. 510/130	2005/0020960	A1 1/2005	Brugger et al.
6,731,971	B2	5/2004	Evans, III et al.	2005/0020961	A1 1/2005	Burbank et al.
6,758,971	B1	7/2004	Haight	2005/0037498	A1 2/2005	Ribi
6,758,975	B2	7/2004	Peabody et al.	2005/0038376	A1 2/2005	Zumeris et al.
6,776,912	B2	8/2004	Baurmeister	2005/0038498	A1 2/2005	Dubrow et al.
6,783,713	B2	8/2004	Tremblay et al.	2005/0048124	A1 3/2005	Sarangapani
6,800,200	B2	10/2004	Bassett et al.	2005/0049181	A1 3/2005	Madhyastha
6,803,363	B2	10/2004	Polaschegg	2005/0059731	A1 3/2005	Albrecht et al.
6,814,085	B2	11/2004	Brattesani et al.	2005/0085775	A1 4/2005	Ostfeld et al.
6,814,724	B2	11/2004	Taylor	2005/0085777	A1 4/2005	Tan
6,835,713	B2	12/2004	Montelaro et al.	2005/0090785	A1 4/2005	Tan
6,838,005	B2	1/2005	Tepper et al.	2005/0095351	A1 5/2005	Zumeris et al.
6,843,784	B2	1/2005	Modak et al.	2005/0100937	A1 5/2005	Holmes
6,844,028	B2	1/2005	Mao et al.	2005/0101841	A9 5/2005	Kaylor et al.
6,849,463	B2	2/2005	Santini, Jr. et al.	2005/0118239	A1 6/2005	Sabesan
6,858,139	B2	2/2005	Taylor	2005/0124678	A1 6/2005	Levy et al.
6,887,270	B2	5/2005	Miller et al.	2005/0131332	A1 6/2005	Kelly et al.
6,887,847	B2	5/2005	Montelaro et al.	2005/0131356	A1 6/2005	Ash et al.
6,921,390	B2	7/2005	Bucay-Couto et al.	2005/0137536	A1 6/2005	Gonnelli
2001/0010016	A1	7/2001	Modak	2005/0142622	A1 6/2005	Sanders et al.
2001/0049975	A1	12/2001	Howard et al.	2005/0143286	A1 6/2005	Singh et al.
2002/0009436	A1	1/2002	Doyle et al.	2005/0154448	A1 7/2005	Cully et al.
2002/0015716	A1*	2/2002	Jampani et al. 424/401	2005/0158253	A1 7/2005	Budny et al.
2002/0037260	A1	3/2002	Budny et al.	2005/0161859	A1 7/2005	Miller et al.
2002/0051812	A1	5/2002	DiCosmo et al.	2005/0171501	A1 8/2005	Kelly
2002/0064858	A1	5/2002	Yacoby-Zeevi	2005/0192590	A1 9/2005	Feeley et al.
2002/0066702	A1	6/2002	Liu	2005/0199245	A1 9/2005	Brennan
2002/0068093	A1	6/2002	Trogolo et al.	2005/0209547	A1 9/2005	Burbank et al.
2002/0091424	A1	7/2002	Biel	2005/0220837	A1 10/2005	Disegi et al.
2002/0102405	A1	8/2002	Chapman et al.	2005/0221072	A1 10/2005	Dubrow et al.
2002/0123787	A1	9/2002	Weiss	2005/0233950	A1 10/2005	Madhyastha
2003/0018306	A1	1/2003	Bucay-Couto et al.	2005/0256502	A1 11/2005	DiMatteo et al.

* cited by examiner

Figure 1



1

INVISIBLE ANTIMICROBIAL GLOVE AND HAND ANTISEPTIC

This application is 371 of PCT/US04/009009, filed 24 Mar. 2004, which claims priority from provisional application No. 60/457,054, filed 24 Mar. 2003.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to an alcohol-based antiseptic surgical scrub, which leaves an anti-microbial invisible film or "glove" on the hands, following alcohol evaporation.

2. Description of Related Art

Diligent hand washing and the wearing of gloves by healthcare professionals are essential components of effective infection control in the healthcare environment. Healthcare professionals regularly wash their hands and wear gloves to control the spread of bacteria and infection from patient to patient, and to themselves.

Hand washing procedures are performed in several ways and include products such as an ordinary anti-microbial bar soap, skin disinfecting or pre-operative agents or rubbing alcohol. Such procedures and products contain anti-microbial agents such as alcohol, iodine, chlorhexidine digluconate, chloroxylenol, triclosan and hexachlorophenes. Gloving by the healthcare workers is a normal practice in the hospital setting to create a barrier that protects themselves and the patient from transmitted bacteria.

Healthcare workers commonly use scrub brushes impregnated with anti-microbial agents for antiseptic hand washing prior to surgical procedures and other invasive patient care procedures. These impregnated scrub brushes have proven to be an effective method of reducing the spread of infection in the healthcare setting. Anti-microbial solutions are specially designed for use in the scrub brush where the mechanical action of scrubbing with the brush creates a foam or lather. Before performing a surgical procedure, the surgical team uses surgical scrubs to disinfect their hands with such a standardized scrub procedure, usually lasting 5 to 10 minutes, then don sterile gloves before initializing the surgical procedure.

Alcohols, in general, are recognized for their disinfection properties. Rubbing alcohol, containing 70% ethyl alcohol (Formula 23-H) and 30% water, and isopropyl rubbing alcohol, containing 70% isopropyl alcohol and 30% water, are commonly used disinfectants. Alcohol is a potent anti-microbial agent, and, if used with surgical scrub, will cause significant mean log reduction of bacterial counts on the hands of healthcare workers. However, the use of alcohols for surgical scrubs has never become popular in the United States of America, even though the compound has very good properties for immediate reduction of microbial flora on the hands. This may be due to the fact that alcohol is a defatting agent. When applied to the human skin, it can make the skin very dry, often leading to chapped and cracked skin. These characteristics thus often result in difficult and painful sensitivity for the users.

The prior art includes many examples of varying antiseptic skin cleaners and wipes, some of which include alcohol. U.S. Pat. No. 4,678,606 discloses a personal liquid cleaning composition having a primary surfactant selected from the group of anionic, cationic, zwitterionic, amphoteric and semipolar surfactants, an auxiliary surfactant selected from certain ethoxylated aliphatic alcohols and a water-soluble polymeric thickening agent. U.S. Pat. No. 4,374,126 discloses an alcohol insoluble antimicrobial topical lotion including a lower acrylate interpolymers. U.S. Pat. No. 5,512,199 discloses an antimicrobial hand wipe which includes an alcohol. U.S. Pat. No. 6,180,584 is directed to a film forming antimicrobial

2

composition. Notably, this composition is an alcohol-free anti-microbial skin sanitizing composition.

However, none of these prior art products provide the benefits of the presently disclosed surgical scrub composition, which not only includes alcohol, but also provides the desired invisible, anti-microbial long-lasting film on the hands following solvent evaporation.

SUMMARY OF THE INVENTION

The present invention is an alcohol based hand surgical scrub, which includes dual anti-microbial agent preservatives, cationic polymer film-forming agents and provides a long term residual, anti-microbial "invisible glove" on the skin. The composition desirably comprises an alcohol, anti-microbial agent which prevents the growth of bacteria and a polymer system to form a barrier on the skin. The polymer system creates an invisible film on the skin following solvent evaporation. This invisible film provides a lasting anti-microbial barrier on the skin.

The composition desirably includes an alcohol, a positive charged anti-microbial agent, skin emollients and lubricants and a film-forming polymer system. The composition may also include one or both of methyl and propyl parabens and silicone. The anti-microbial agents prevent bacteria from growing back to the baseline of the normal skin flora population. The surgical scrub composition contains chemical ingredients to remove extraneous matter and organisms from the hands. Skin emollients and lubricants are also provided to prevent the defatting effect of the alcohol, thereby eliminating the dry, chapped feeling on the skin. Additional or complementary anti-microbial agents are preferably included to serve as preservative and persistent active agents. These anti-microbial agents in the composition will improve the antimicrobial characteristics of the formulation and provide a long-term residual activity for the skin. As previously noted, the polymer system is included to create an invisible film after the solvent is evaporated. This invisible film acts as a physical anti-microbial barrier or "glove" to protect the healthcare worker's hands. This film has an anti-microbial effect at least for six hours to prevent and kill the bacteria on the healthcare worker's hands.

DESCRIPTION OF THE PREFERRED EMBODIMENT(S)

This invention may be satisfied by embodiments in many different forms, and is not limited to any specific or preferred embodiments of the invention, which are merely exemplary of the principles of the invention. Various other modifications will be apparent to and readily made by those skilled in the art without departing from the scope and spirit of the invention. The scope of the invention will be measured by the appended claims and their equivalents.

The surgical scrub composition of the present invention includes alcohol, a positive-charged anti-microbial agent, skin emollients and lubricants, and a cationic polymer film-forming thickening agent.

One or more alcohols are utilized in the surgical scrub composition due to alcohol's well known germicidal properties. Alcohols which may be used in the present composition include isopropyl alcohol, ethanol, and n-propanol. In the most preferred embodiment, ethyl alcohol is used. Preferably, alcohol is present in the surgical scrub composition in an amount from about 60 to about 95 weight percent, and most preferably, at about 71 weight percent.

Emollients are utilized in the surgical scrub composition of the present invention to lessen the drying effect of the alcohol on the skin. A preferred emollient is isopropyl palmitate, which is an ester emollient available from Amerchol Corpo-

ration of Edison, N.J. When used on the skin, isopropyl palmitate leaves a soft, non-sticky feel to the skin. Isopropyl palmitate is a lipophilic emollient, an oily, fatty substance which is water insoluble. Lanolin and its derivatives are a large group of fatty materials with active emollient properties that can be utilized as emulsifiers or lubricants in the composition.

The preferred embodiment includes glycerin as a hydrophilic emollient and a humectant to maintain the water in equilibrium. Glycerin is available from Spectrum Quality products. Glycerin is a water and alcohol soluble emollient that provides skin conditioning properties to the composition. In another embodiment, propylene glycol may be substituted for glycerin. Preferably, glycerin is present in the surgical scrub composition, by weight percent of the entire composition, in an amount of from about 0.1% to about 5%, and most preferably, about 1.4%.

The surgical scrub composition further includes a silicone fluid. Preferably, the silicone fluid is dimethicone copolyol, cyclomethicone and dimethicone. Most preferably, dimethicone is utilized. Dimethicone 350 is commercially available from General Electric Corporation. Silicone fluids provide good lubricating, aesthetic and occlusive action on the skin. Silicone fluids not only lubricate the skin surface, they also act as a gas barrier to reduce water vapor transmission. Silicone fluids are water and gas insoluble. Preferably, the silicone fluid is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from 0 to about 5%, and most preferably, about 0.03%.

Most hydrocarbons are derived from petrolatum crude by a fractional distillation process. Such hydrocarbons function as emollients due to their ability to lubricate and hold water at the skin surface due to their relative occlusivity. Mineral oil and isopropyl palmitate are included in the most preferred embodiment. Despite its lack of aesthetic properties, petrolatum may be utilized. Alternatively or additionally, lanolin derivatives may be included. Preferably, mineral oil is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from 0 to about 5%, and most preferably, about 0.2%. Preferably, isopropyl palmitate is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0.1% to about 5%, and most preferably about 0.4%.

Preferably, the surgical scrub composition includes about 0.1% to about 5% of an emollient or emollients.

An anti-microbial agent is included in the surgical scrub composition to kill microorganisms and prevent or inhibit their growth and reproduction. In the absence of an antimicrobial agent, microbial flora will grow on the skin following alcohol evaporation. The antimicrobial agents present in the surgical scrub composition are selected to be compatible with the chemical and physical properties of the skin. In addition, the antimicrobial agent should be non-toxic and environmentally friendly.

As previously noted, the alcohol component of the surgical scrub composition is an excellent antimicrobial agent. It also acts as a preservative. However, following application of the surgical scrub composition, the alcohol will evaporate from the skin over time. In the most preferred embodiment, the residue remaining on the skin following alcohol evaporation will include an amount of an antimicrobial agent which also acts as a preservative. Suitable antimicrobial agents which act as preservatives may be selected from the class of phenols including para-chloro-meta-xyleneol, bis-biguanides such as chlorhexidine digluconate, chlorhexidine diacetate or quaternium class such as benzethonium chloride, benzalkonium chloride. Chloroxylenol, triclosan, hexachlorophenes, octenidine and quarternary compounds may be included. Hexetidine, germaben II, kathon CG, triclosan are other antimicrobial agents may also be suitable as preservatives. Benzethonium chloride and benzalkonium chloride (as hyamine

3500) are available from Lonza Inc., chlorhexidine digluconate is available from Xtrium Laboratories, germaben II is available from Sutton Laboratories. In a most preferred embodiment, benzethonium chloride, chlorhexidine digluconate (20%) and benzalkonium chloride (80%) are present. Preferably, benzethonium chloride is present in the scrub composition, by weight percent of the entire composition, in an amount from about 0.05% to about 5%, and most preferably, about 0.09%. Preferably, chlorhexidine digluconate (20%) is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0.05% to about 5%, and most preferably about 0.45%. Preferably, benzalkonium chloride (80%) is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from 0 to about 5%, and most preferably, about 0.09%. Preferably, the composition includes about 0% to about 5% of an anti-microbial agent.

Methylparaben and propylparaben (both available from Mallinckrodt Chemical Company) are desirably included in the surgical scrub composition as preservatives. These are further desired for their moisturizing and film forming characteristics. Preferably, methylparaben is contained in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0 to about 5%, and most preferably about 0.03%. Preferably, propylparaben is present in the surgical scrub, by weight percent of the entire composition, in an amount from about 0 to about 5%, and most preferably, about 0.03%. One or both of these compositions are preferably included in the amount of about 0% to about 5% of the surgical scrub composition.

To create the desired residual film of the surgical scrub composition, cationic polymer thickening agents are employed. Thickening agents must be soluble in the alcohol and compatible with cationic ingredients. The cationic polymer thickening agents are preferably present in the surgical scrub composition, by weight percent of the entire composition, in the amount of about 0.1% to about 5%.

Incroquat behenyl TMS is most preferably used as a cationic polymer. It is a strong conditioning agent and an outstanding cationic emulsifier. This composition bonds to the skin due to the skin's negative charge. Incroquat behenyl TMS is a compound of 25% active solution of behenyl trimonium methosulfate in cetearyl alcohol and is available in flaked/pastel form and supplied by Croda Inc. Incroquat CR concentrate is preferably used as an additional cationic polymer. Incroquat CR concentrate is available from Croda Inc. and is comprised of cetearyl alcohol, castor oil and stearylalkonium chloride. This composition is a one-part formulating aid. It is a conditioner and self-emulsifier. The desired combination of the incroquat behenyl TMS and incroquat CR concentrate provides a smooth feel to the skin and neutralizes the static charge of the human skin. Preferably, incroquat BTMS is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0.1% to about 5%, and most preferably about 0.4%. Preferably, incroquat CR is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from 0 to about 5%, and most preferably about 0.1%.

Alternative or additional polymers include cellulose, polyvinylpyrrolidone and waxes.

The most preferred embodiment of the surgical scrub composition is as follows:

	% of Total (w/w)
1-Docosanol	1.00%
Benzethonium Chloride	0.09%
Chlorhexidine Digluconate, (20%)	0.45%
Dimethicone	0.03%

-continued

	% of Total (w/w)
Ethyl Alcohol	71.00%
Glycerin	1.40%
Benzalkonium Chloride	0.09%
Incroquat BTMS	0.40%
Incroquat CR	0.10%
Isopropyl Palmitate	0.40%
Mineral Oil	0.20%
Methylparaben	0.03%
PEG-10 Behenyl Ether	1.50%
Propylparaben	0.03%
Purified Water	23.28%

The most preferred embodiment includes the additional components 1-Docosanol as a moisturizer and film forming substance, PEG-10 behenyl ether as a moisturizer, fragrance and water. Preferably, 1-Docosanol is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0.1% to about 5%, and most preferably about 1%. Preferably, PEG-10 behenyl ether is present in the surgical scrub composition, by weight percent of the entire composition, in an amount from about 0.1% to about 5%, and most preferably about 1.5%. Preferably, purified water is present in the surgical scrub composition, in an amount from about 1% to about 35%, and most preferably about 23%.

The surgical scrub composition does not cause skin irritation or sensitization. Additional advantages of the composition include its compatibility with positive-charged anti-mi-

The following examples are not intended to limit the scope of the invention, but are only intended to be exemplary in nature.

Standard testing of the most preferred embodiment demonstrates its effectiveness as an antiseptic to provide an immediate reduction of microbes and then to prevent microbial growth back to baseline within a 6-hour period. The immediate log reduction was 2.65 on day 1, 2.92 on day 2 and 3.04 on day 3. The residual effect was a log reduction six hours after hand cleaning of 1.45 on day 1, 1.79 on day 2 and 2.18 on day 3. The residual anti-microbial activity is facilitated by the containment of the anti-microbial agent in the polymer film that is formed. The film forms an invisible "glove" containing an anti-microbial agent. The anti-microbial agent remains on the skin in the film and continues to be bactericidal over extended periods of time. The invisible glove prevents microbes that may be present on the hands of the healthcare worker from infecting a patient and the glove also provides a barrier for microbes infecting the healthcare worker from the patient.

Through testing, the composition has been shown to exceed industry standards. The critical indices for this study were an immediate one (1) \log_{10} reduction in microorganisms on Day 1; an immediate two (2) \log_{10} reduction in microorganisms on Day 2; an immediate three (3) \log_{10} reduction in microorganisms on Day 5; and that microorganism populations from the approximately three (3) hour to three (3) hour and thirty (30) minute AND approximately six (6) hour to six (6) hour and thirty (30) minute samples not return to baseline level. Refer to Table I.

TABLE I

Statistical Summary of the \log_{10} Recovery Values of Most Preferred Embodiment						
Sample	Sample Size	Mean	Standard Deviation	95% Confidence Interval	\log_{10} Reduction from Baseline	TFM (1994) Required \log_{10} Reduction
Baseline	36	6.01	0.40	5.88 to 6.15	N/A	
Day 1, Immediate	12	3.36	0.68	2.93 to 3.79	2.65	1
Day 1, 3 Hour	12	4.08	0.99	3.45 to 4.71	1.93	No return to baseline
Day 1, 6 Hour	12	4.56	0.68	4.13 to 4.99	1.45	No return to baseline
Day 2, Immediate	11	3.09	0.53	2.74 to 3.45	2.92	2
Day 2, 3 Hour	11	3.73	0.90	3.12 to 4.34	2.28	No return to baseline
Day 2, 6 Hour	12	4.22	0.81	3.70 to 4.73	1.79	No return to baseline
Day 5, Immediate	10	2.97	0.47	2.63 to 3.31	3.04	3
Day 5, 3 Hour	11	3.50	0.61	3.09 to 3.91	2.51	No return to baseline
Day 5, 6 Hour	11	3.83	1.01	3.15 to 4.51	2.18	No return to baseline

crobial agents such as chlorhexidine digluconate, thus ensuring the effectiveness of the anti-microbial agent or agents. The included skin emollients and lubricants eliminate skin drying or chapping. These compositions counteract the defatting effect of the alcohol.

The surgical scrub composition includes anti-microbial agents, which act as both persistent active agents and suitable preservatives. This composition permits a very small amount of an anti-microbial agent to improve anti-microbial activities and provide long-term residual activity for the skin.

Traditionally, healthcare workers have scrubbed prior to every surgical procedure. An improved method may be to apply water to the skin, perform hand scrubbing at the beginning of the day using a scrub brush and anti-microbial scrub solution, rinsing the anti-microbial scrub solution with water, and then utilization of the composition of the present invention prior to subsequent surgical procedures. A high level of hand antisepsis can be achieved via this method without abrasive scrubbing multiple times a day.

The most preferred embodiment was also tested in conjunction with a standard hand scrub utilizing a brush impregnated with an anti-microbial agent. The composition was applied to the hands of the test subjects one-half hour following the standard scrub. Test results showed an increased log reduction of microbes when the scrubbing and the preferred composition were used together. Log reductions of microbes were 3.66 for Day 1, 3.98 for Day 2, and 3.81 for Day 3. Residual activity was also increased with log reductions after six hours of 2.31 for Day 1, 3.52 for Day 2 and 3.70 for Day 3. Use of the preferred composition in conjunction with a standard scrub exceeded industry standards for anti-microbial compositions. Refer to Table II and FIG. 1.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graphical presentation of the \log_{10} reductions from baseline for the preferred formula and BD E-Z Scrub™ [BD Catalog #371073] plus the most preferred embodiment.

TABLE II

Statistical Summary of the Log ₁₀ Recovery Values Using BD E-Z Scrub™ Catalog #371073 Plus Most Preferred Embodiment						
Sample	Sample Size	Mean	Standard Deviation	95% Confidence Interval	Log ₁₀ Reduction from Baseline	TFM (1994) Required Log ₁₀ Reduction
Baseline	38	6.10	0.38	5.97 to 6.22	N/A	
Day 1, Immediate	12	2.88	0.46	2.15 to 2.73	3.66	1
Day 1, 3 Hour	13	3.13	0.79	2.65 to 3.60	2.97	No return to baseline
Day 1, 6 Hour	13	3.79	0.88	3.26 to 4.32	2.31	
Day 2, Immediate	12	2.12	0.35	1.90 to 2.34	3.98	2
Day 2, 3 Hour	12	2.58	0.93	1.99 to 3.17	3.52	No return to baseline
Day 2, 6 Hour	12	2.58	0.66	2.16 to 3.00	3.52	
Day 5, Immediate	10	2.29	0.37	2.02 to 2.55	3.81	3
Day 5, 3 Hour	10	2.28	0.32	2.05 to 2.51	3.82	No return to baseline
Day 5, 6 Hour	12	2.40	0.49	2.09 to 2.71	3.70	

The invention claimed is:

1. A surgical scrub composition comprising, in weight percent of the total composition:

- (a) from about 0.1% to about 5% of 1-docosanol;
- (b) from about 0.05% to about 5% of chlorhexidine digluconate;
- (c) from about 60% to about 95% of alcohol;
- (d) from about 0.1% to about 5% of glycerin;
- (e) from about 0.1% to about 5% of a 25% active solution of behenyl trimethyl ammonium methosulfate in cet-
earyl alcohol;
- (f) from about 0.1% to about 5% of isopropyl palmitate;

(g) from about 0.03% to about 5% of a preservative selected from the group consisting of methylparaben and propylparaben; and

(h) from about 1% to about 35% of purified water.

2. A method of using a surgical scrub composition for hand antisepsis comprising:

- (a) applying a cleansing liquid to the skin;
- (b) scrubbing the skin with an anti-microbial scrub solution;
- (c) rinsing the anti-microbial scrub solution from the skin; and
- (d) applying an effective amount of a surgical scrub composition according to claim 1.

* * * * *